AD/A-000 534

ON THE QUESTION OF SYNCHRONISM OF CLIMATIC CHANGES IN THE NORTH ATLANTIC-ARCTIC AND NORTH PACIFIC ARCTIC

I. I. Schell, et al

Ocean-Atmosphere Research Institute

Prepared for:

Air Force Office of Scientific Research Advanced Research Projects Agency

August 1974

DISTRIBUTED BY:



REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM		
AFOSR - TE + 74 - 1612	3. RECIPIENT'S CATALOG NUMBER		
On The Question of Synchronism of Climatic Changes	5. Type of REPORT & PERIOD COVERED Final Report June 1, 1972-June 30.1		
in the No. Atlantic-Arctic and No. Pacific Arctic	6. PERFORMING ORG. REPORT NUMBER		
I. I. Schell and E. N. Sabbagh	8. CONTRACT OR GRANT NUMBER(3) F144620-72-C-0040		
Ocean-Atmosphere Research Institute 1640 Massachusetts Avenue Cambridge, Lass. 02138	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS AO 2159 62701D		
I. CONTROLLING OFFICE NAME AND ADDRESS Advanced Research Projects Agency	2. REPORT DATE AUGUST, 1974		
1400 Wilson Boulevard Arlington, VA 22209	13. NUMBER OF PAGES		
Air Force Office of Scientific Research /NP	Unclassified		
1400 Wilson Boulevard Arlington, VA 22209			

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

None

19. KEY WORDS Continue on reverse side if necessary and identify by block number) Climatic Changes

Synchronism

No. Atlantic - Arctic

No. Pacific - Arctic

20. ABSTRACT (Continue on severse side if necessary and identify by block number)
An analysis of the climatic changes from the 1921-50 period to the following 1951-60 and 1961-70 decades showed moderate synchronism in the annual and December-Way changes of the circulation of the two regions as revealed by the intensity and position of the No. Pacific (Aleutian) and No. Atlantic (Icelandic) Lows. A filling-in and southwestward retreat of the Aleutian Low in the 1950's and 1960's in relation to the 1921-50 period was parallele! by a filling-in of the Icelandic Low in the same periods.

DD 1 JAN 73 1473

TION OF 1 NOV 65 IS OBSOLETE

Unclassified

li

and a further lowering in the 1960's in Alaska and Iceland and also a lowering of the December-May temperatures in both regions in the 1950's and a further lowering in the 1960's in Iceland though not in Alaska, possibly due to a change in the site of the stations in the 1960's.

Ocean-Atmosphere Research Institute 1640 Massachusetts Avenue Cambridge, Massachusetts 02138

AFOSR - TR - 74 - 1612


FINAL REPORT

I. I. Schell and E. N. Sabbagh

August 1974

Sponsored by
Advanced Research Projects Agency
ARPA Order No. 2159

Approved for public release; distribution unlimited.



ARPA Order No. 2159
Program Code - F10
Name of Contractor - Ocean-Atmosphere Research Institute
Effective Date of Contract - June 1, 1972
Contract Expiration Date - June 30, 1974
Amount of Contract Dollars - \$\pi\$104,007
Contract Number - F44620-72-C-0040
Principal Investigator and Program Manager - I. I. Schell
Phone Number - 617-491-8552
Short Title of Work: Synchronism of Climatic Changes

NATIONAL TECHNICAL INFORMATION SERVICE

The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency of the U.S. Government.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO THE
THE ASSOCIATION OF THE COLOR AND THE 150-12 (7b).
DESCRIPTION OF THE COLOR AND THE 150-12 (7b).
D. W. TAYLOR

Technical Information Officer

Summary Report

An investigation was made of the degree of synchronism of climatic changes in the No. Atlantic-Arctic and No. Pacific-Arctic from a consideration of the changes in the circulation (pressure distribution), temperature, and ice conditions in Bering Strait and off Iceland to help us obtain some ideas of the causes of the climatic changes. An increased amount of heat, for example, reaching the surface of the earth due to a possible increase in the sun's heat output or to a decrease in the atmospheric turbidity for different causes would in time call for a global warming or a synchronous rise in temperature in all regions while an internal adjustment of the atmospheric circulation involving also the oceans would lead to a warming in some regions and cooling in others or a non-synchronous change.

An analysis of the changes in the circulation from the 1921-50 period to the following 1951-60 and 1961-70 decades showed moderate synchronism in the annual and December-May changes of the circulation of the two regions as revealed by the intensity and position of the No. Pacific (Aleutian) and No. Atlantic (Icelandic) Lows. Thus, the filling-in and south-westward retreat of the Aleutian Low in the 1950's and 1960's in relation to the 1921-50 period was paralleled by a filling-in of the Icelandic Low in the same periods together with a related southward intrusion of the Arctic High towards Iceland. This was accompanied by a lowering of the annual temperatures

in the 1950's and a further lowering in the 1960's in Alaska and Iceland and also a lowering of the December-May temperatures in both regions in the 1950's and a further lowering in the 1960's in Iceland through not in Nome and Bethel, possibly due to a change in the site of the stations in the 1960's.

The sharp increase in the ice off Iceland in the 1960's presumably had its origin in the increased transport of the ice from the pack in the North that manifested itself first north of Iceland in the 1950's.

On the Question of Synchronism of Climatic Changes in the No. Pacific and No. Atlantic Arctic

By I. I. Schell and E. N. Sabbagh Ucean-Atmosphere Research Institute Cambridge, Mass. 02138

A. Introduction

An important consideration in studies of possible causes of climatic changes is the degree of synchronism of the changes of different regions; whether or not they develop more or less simultaneously and in the same direction or not in various regions depending on the cause of the climatic change. An increased amount of heat, for example, reaching the surface of the earth due to a possible increase in the sun's heat output or to a decrease in the atmospheric turbidity with a decrease in volcanism would in time call for a global warming or a synchronous rise in temperature in all regions while internal adjustments of the atmospheric circulation involving also the oceans would lead to a warming in some regions and a cooling in others or a non-synchronous change.

Atlantic-Arctic can differ markedly in a given year from those in the North Pacific-Arctic was recently presented by Rodewald (1958) by comparing phenomenally heavy ice conditions off northern Alaska in 1955 with the light ice conditions in the

however, that over a period measured in decades, rather than in years, the response to a cause of climatic change would show itself as a trend. To learn whether changes in ice conditions of the order of a decade or more in the North Atlantic-Arctic are relatively synchronous with those in the North Pacifie-Arctic, a study was made of the ice conditions off Iceland and in Bering Strait along with the circulation (pressure distribution) and air temperatures of the three different periods 1921-50, 1951-60, 1961-70 and compared for the two regions.

- B. Ice off Iceland and in Bering Strait
- is usually expressed by an index that measures the duration and extent of the ice along the coast. The average value of this index for the 30-year period (1921-50) was 3.1 units followed by 1.1 units in the 1951-60 decade and by a sharp rise in the last decade (1961-69) that came to 50.4 units (Eythorsson and Sigtryggsson, 1971).
- Dates. In the absence of historical data on the ice limit in the Bering Sea, we turn to the long record of opening and closing dates to navigation of St. Fichael's in Norton Sound, opening on Bering Strait (Weightman, 1941). While the dates of opening and closing in individual years at a given location are not likely to reflect ice conditions over a wide area, we may assume that over a longer period, one or more decades, it

may. The average duration of the ice season at St. Michael for the 1921-50 period is 206 days followed by 202 days in the 1950's, while the dates for the 1960's are still being processed by the National Climatic Center in Asheville (see also Weightman, 1941).

C. No. Atlantic (Icelandic) Low

1. Annual. Fig. 1 giving the average annual pressure distribution in the No. Atlantic-Arctic shows isobar 1008 mb north of Icoland to run westward to the coast towards Greenland indicating little ice movement towards Iceland. A movement of ice towards Iceland in the 1950's is indicated (Fig. 2) from the direction of isobar 1012 mb which runs southwestward and a still greater movement in the 1960's (Fig. 3) as indicated south from this isobar which runs/southwestward towards the coast.

Along with the indicated increased southward movement of the ice from the pack in the North in the 1950's and a still greater movement in the 1960's, we note a lowering in temperatures at Stykkisholm and Teigarhorn, respectively, in Iceland in the 1950's and an additional lowering in the 1960's as the cold air from the north penetrates farther southward. This is revealed also from a filling-in of the Icelandic Low in the 1950's as shown by a shrinkage of the area enclosed by the 1008 mb isobar and a further filling-in of the Low in the 1960's shown by a further reduction of the enclosed area (Figs.1-3) as well as the disappearance of the area enclosed by the 1006 mb isobar in that decade.




Fig. 1 1921-1950 Annual Pressure (mb).




Fig. 2 1951-1960 Annual Pressure (mb).




Fig. 3 1961-1970 Annual Pressure (mb).

2. <u>Dec.-May</u>. A similar filling-in of the Icelandic Low may be seen in the 1950's and another filling-in in the 1960's during the December-Lay season as shown by the shrinking of the area enclosed by the 1004 mb isobar in the 1950/1-1959/60 period (Fig. 5) as compared with the 1920/1-1949/50 period (Fig. 4) and its disappearance in the 1960/1-1969/70 (Fig. 6.). This is associated with a southward extension of the Arctic High (1016 mb isobar) in the 1950's (Fig. 5) from its position in the preceding 30-year period (Fig. 4) and a further southward extension of the High in the 1960's (Fig. 6), together indicating a progressively greater outflow of cold air and a southward movement of the ice from the pack in the North towards Iceland that finally reaches the coast in the 1960's.

D. No. Pacific-Arctic (Aleutian Low)

1. Annual. Fig. 7 shows the Aleutian Low in 1921-50 to be well developed with isobar 1008 mb extending northeastward into southwestern Alaska in contrast to the same isobar extending only into the Alaskan Peninsula in 1951-60 (Fig. 8) and even less northeastward stopping short of the Peninsula in the 1961-70 decade (Fig. 9), reflecting a progressive southwestward withdrawal of the Low in the 1951-60 and 1960-70 decades. The southwestward withdrawal of the Aleutian Low in 1951-60 and again in 1961-70 suggests a lesser inflow of relatively warm air into the region in the last two decades that would account for the lowering of the annual temperature at Nome and Bethel in 1951-60 as compared with the 1921-50 period and a further lowering in 1961-70 (see Table 1).




Fig. 4. December-May 1921-1950 Pressure (mb).




Fig. 5. December-Nay 1951-1960 Pressure (mb).




Fig. 6. December-May 1961-1970 Pressure (mb).




Fig. 7. 1921-1950 Annual Pressure (mb).




Fig. 8. 1951-1960 Annual Pressure (mb).




Fig. 9. 1961-1970 Annual Pressure (mb).

2. <u>December-May</u>. A similar trend appears in the December-May season when St. Michael is on the average beset by ice as reckoned in whole months. Figs. 10-12 show isobar 1008 mb reaching northeastward into southwestern Alaska in 1921-50, and again stopping short of the Peninsula in 1961-70, showing a progressive withdrawal of the Low southwestward. A filling-in of the Aleutian Low in the 1951-60 and 1961-70 periods is revealed from the absence of the 1004 mb isoabar shown in the 1921-50 period (Fig. 10). This is accompanied by lowering temperatures in both the 1950's and the 1960's (see below).

E. Temperature

Table 1 giving the mean annual as well as December-May temperatures at Stykkisholm and Teigarhorn (Iceland) shows a decrease in the 1951-60 decade from its 1921-50 averages and a further decrease in the 1961-70 decade, in accordance with an increased inflow of cold air from the north in the 1950's and again in the 1960's as compared with the 1921-50 period.

Table 1 similarly giving the mean annual as well as the December-May temperatures at the Alaskan stations Nome and Bethel opening on sering Strait shows a decrease during the 1951-60 decade from the average of the preceding 1921-50 period and a further decrease in the annual though not in December-way, in the 1961-70 decade (see below), in keeping with the mouthwestward withdrawal of the Aleutian Low and a presumed leaper inflow of relatively warm air from the ocean modified by its passage over land.






Fig. 10.December-May 1920/1-1949/50 Pressure (mb).






Fig. 11. December-May 1950/1-1959/60 Pressure (mb).




Fig. 12. December-May 1960/1-1969/70 Pressure (mb).

We are at a loss to explain the lack of a further decrease in the December-May temperatures at Bethel and Nome in the 1960's unless it arose from a change in the site at Bethel and to an elevation of the thermometer to 125 feet from 10 feet, which would raise its temperature in winter due to the inversion, and also a move of the Nome station from its previous site.

Table 1. Mean annual and December-May temperature deviations at Nome and Bethel in Alaska and at Stykkisholm and Teigarhorn in Iceland in the 1951-60 and 1961-70 decades from their respective 1921-50 averages.

	Nome		Bethel		Stykkisholm		Teigarhorn	
	Ann.	Dec-May	Ann.	Dec-May	Ann.	Dec-Lay	Ann.	Dec-hay
	oc	oc	oc	oC	oC	оС	oc	oc
1951-60	-0.2	-0.9	-0.5	-1.3	-0.1	-0.3	-0.2	-0.6
1 961 - 70	-0.6	-0.6	-1.2	-0.8	-0.7	-0.7	-0.8	-0.8
1921-50 (average)	-3.2	-10.2	-1.4	-8.3	4.1	1.4	4.6	2.3

F. Remarks

Our attempt to show synchronism in the changes in the No. Atlantic-Arctic and the No. Pacific Arctic by comparing the circulation (pressure distribution), air temperatures, and the ice conditions off Iceland with those in Bering Strait can be said to have been moderately successful.

A moderate degree of synchronism between the two regions is shown by the southwestward retreat and filling-in of the Aleutian Low in the 1951-60 decade as compared with the 1921-50 period preceding it, and again a further retreat and a filling-in

ressive shrinking and a filling-in of the No. Atlantic Low and a southward extension of the Arctic High over northern Greenland affecting Iceland in the 1950's and 1960's on the other. Similarly, the lowering of the annual and December-Lay temperatures in the 1950's in both regions and the further lowering of the annual and December-Lay temperatures in the 1950's in Iceland and possibly also in Alaska signifies again a certain synchronism in the climatic changes of the two regions.

The record of opening and closing dates to navigation at St. Michael opening up on Bering Strait in the 1960's, as this is being written, is not yet available to allow a comparison with the ice conditions off Iceland.

Acknowledgments: We are indebted to the officials of the U.S. Mational Climatic Data Center, NOAA, and the U.S. Naval Oceanomraphic Office for kindly furnishing ice data on St. Michael used in this report.

References

- Eythorsson, J. and H. Sigtryggsson 1971: The Climates and Weather of Iceland. In "Zoology of Iceland.." 1(3): 1-62.

 E. Munksgaard (Copenhagen and Reykjavik)
- Rode vald, M. 1968: Die "Eis-Schaukel" im nordlichen Polarmeer im Jahre 1955. Ann. Meteor. 8(5/6): 187-9.
- Forecasting. M. Wea. Rev. Suppl. No. 45. 99 Pp.